

Ma

KEY STAGE

3

ALL TIERS

2005

Mathematics tests

Mark scheme for  
Paper 2

Tiers 3–5, 4–6, 5–7 and 6–8

2005



department for

**education and skills**

creating opportunity, releasing potential, achieving excellence

# Introduction

The test papers will be marked by external markers. The markers will follow the mark scheme in this booklet, which is provided here to inform teachers.

This booklet contains the mark scheme for paper 2 at all tiers. The paper 1 mark scheme is printed in a separate booklet. Questions have been given names so that each one has a unique identifier irrespective of tier.

## The structure of the mark schemes

The marking information for questions is set out in the form of tables, which start on page 11 of this booklet. The columns on the left-hand side of each table provide a quick reference to the tier, question number, question part, and the total number of marks available for that question part.

The **Correct response** column usually includes two types of information:

- a statement of the requirements for the award of each mark, with an indication of whether credit can be given for correct working, and whether the marks are independent or cumulative;
- examples of some different types of correct response, including the most common.

The **Additional guidance** column indicates alternative acceptable responses, and provides details of specific types of response that are unacceptable. Other guidance, such as when 'follow through' is allowed, is provided as necessary.

Questions with a *UAM* element are identified in the mark scheme by an encircled *U* with a number that indicates the significance of using and applying mathematics in answering the question. The *U* number can be any whole number from 1 to the number of marks in the question.

For graphical and diagrammatic responses, including those in which judgements on accuracy are required, marking overlays have been provided at the centre page of this booklet.

The 2005 key stage 3 mathematics tests and mark schemes were developed by the Mathematics Test Development Team at QCA.

# General guidance

## Using the mark schemes

Answers that are numerically equivalent or algebraically equivalent are acceptable unless the mark scheme states otherwise.

In order to ensure consistency of marking, the most frequent procedural queries are listed on the following two pages with the prescribed correct action. This is followed by further guidance relating to marking of questions that involve money, time, algebra, coordinates, negative numbers or probability. Unless otherwise specified in the mark scheme, markers should apply the following guidelines in all cases.

**What if ...**

<i>The pupil's response does not match closely any of the examples given.</i>	Markers should use their judgement in deciding whether the response corresponds with the statement of requirements given in the <b>Correct response</b> column. Refer also to the <b>Additional guidance</b> .
<i>The pupil has responded in a non-standard way.</i>	Calculations, formulae and written responses do not have to be set out in any particular format. Pupils may provide evidence in any form as long as its meaning can be understood. Diagrams, symbols or words are acceptable for explanations or for indicating a response. Any correct method of setting out working, however idiosyncratic, is acceptable. Provided there is no ambiguity, condone the continental practice of using a comma for a decimal point.
<i>The pupil has made a conceptual error.</i>	In some questions, a method mark is available provided the pupil has made a computational, rather than conceptual, error. A computational error is a slip such as writing $4 \times 6 = 18$ in an otherwise correct long multiplication. A conceptual error is a more serious misunderstanding of the relevant mathematics; when such an error is seen no method marks may be awarded. Examples of conceptual errors are: misunderstanding of place value, such as multiplying by 2 rather than 20 when calculating $35 \times 27$ ; subtracting the smaller value from the larger in calculations such as $45 - 26$ to give the answer 21; incorrect signs when working with negative numbers.
<i>The pupil's accuracy is marginal according to the overlay provided.</i>	Overlays can never be 100% accurate. However, provided the answer is within, or touches, the boundaries given, the mark(s) should be awarded.
<i>The pupil's answer correctly follows through from earlier incorrect work.</i>	Follow through marks may be awarded only when specifically stated in the mark scheme, but should not be allowed if the difficulty level of the question has been lowered. Either the correct response or an acceptable follow through response should be marked as correct.
<i>There appears to be a misreading affecting the working.</i>	This is when the pupil misreads the information given in the question and uses different information. If the original intention or difficulty level of the question is not reduced, deduct one mark only. If the original intention or difficulty level is reduced, do not award any marks for the question part.
<i>The correct answer is in the wrong place.</i>	Where a pupil has shown understanding of the question, the mark(s) should be given. In particular, where a word or number response is expected, a pupil may meet the requirement by annotating a graph or labelling a diagram elsewhere in the question.

**What if ...**

<i>The final answer is wrong but the correct answer is shown in the working.</i>	Where appropriate, detailed guidance will be given in the mark scheme and must be adhered to. If no guidance is given, markers will need to examine each case to decide whether:	
	the incorrect answer is due to a transcription error;	If so, award the mark.
	in questions not testing accuracy, the correct answer has been given but then rounded or truncated;	If so, award the mark.
	the pupil has continued to give redundant extra working which does not contradict work already done;	If so, award the mark.
	the pupil has continued, in the same part of the question, to give redundant extra working which does contradict work already done.	If so, do not award the mark. Where a question part carries more than one mark, only the final mark should be withheld.
<i>The pupil's answer is correct but the wrong working is seen.</i>	A correct response should always be marked as correct unless the mark scheme states otherwise.	
<i>The correct response has been crossed or rubbed out and not replaced.</i>	Mark, according to the mark scheme, any legible crossed or rubbed out work that has not been replaced.	
<i>More than one answer is given.</i>	If all answers given are correct or a range of answers is given, all of which are correct, the mark should be awarded unless prohibited by the mark scheme. If both correct and incorrect responses are given, no mark should be awarded.	
<i>The answer is correct but, in a later part of the question, the pupil has contradicted this response.</i>	A mark given for one part should not be disallowed for working or answers given in a different part, unless the mark scheme specifically states otherwise.	

**Marking specific types of question**

<b>Responses involving money</b> <i>For example: £3.20 £7</i>	
<b>Accept ✓</b>	<b>Do not accept ✗</b>
<p>✓ Any unambiguous indication of the correct amount eg £3.20(p), £3 20, £3,20, 3 pounds 20, £3-20, £3 20 pence, £3:20, £7.00</p> <p>✓ The £ sign is usually already printed in the answer space. Where the pupil writes an answer other than in the answer space, or crosses out the £ sign, accept an answer with correct units in pounds and/or pence eg 320p, 700p</p>	<p>✗ Incorrect or ambiguous use of pounds or pence eg £320, £320p or £700p, or 3.20 or 3.20p not in the answer space.</p> <p>✗ Incorrect placement of decimal points, spaces, etc or incorrect use or omission of 0 eg £3.2, £3 200, £32 0, £3-2-0, £7.0</p>

<b>Responses involving time</b> <i>A time interval For example: 2 hours 30 mins</i>	
<b>Accept ✓</b>	<b>Take care ! Do not accept ✗</b>
<p>✓ Any unambiguous indication eg 2.5 (hours), 2h 30</p> <p>✓ Digital electronic time ie 2:30</p>	<p>✗ Incorrect or ambiguous time interval eg 2.3(h), 2.30, 2-30, 2h 3, 2.30min</p> <p>! The time unit, hours or minutes, is usually printed in the answer space. Where the pupil writes an answer other than in the answer space, or crosses out the given unit, accept an answer with correct units in hours or minutes, unless the question has asked for a specific unit to be used.</p>
<b>A specific time For example: 8.40am, 17:20</b>	
<b>Accept ✓</b>	<b>Do not accept ✗</b>
<p>✓ Any unambiguous, correct indication eg 08.40, 8.40, 8:40, 0840, 8 40, 8-40, twenty to nine, 8,40</p> <p>✓ Unambiguous change to 12 or 24 hour clock eg 17:20 as 5:20pm, 17:20pm</p>	<p>✗ Incorrect time eg 8.4am, 8.40pm</p> <p>✗ Incorrect placement of separators, spaces, etc or incorrect use or omission of 0 eg 840, 8:4:0, 084, 84</p>



<b>Responses involving the use of algebra</b>	
<i>For example: <math>2 + n</math> <math>n + 2</math> <math>2n</math> <math>\frac{n}{2}</math> <math>n^2</math></i>	
<b>Accept ✓</b>	<b>Take care ! Do not accept ✗</b>
<p>✓ Unambiguous use of a different case or variable eg <math>N</math> used for <math>n</math> <math>x</math> used for <math>n</math></p>	<p>! Unconventional notation eg <math>n \times 2</math> or <math>2 \times n</math> or <math>n2</math> or <math>n + n</math> for <math>2n</math> <math>n \times n</math> for <math>n^2</math> <math>n \div 2</math> for <math>\frac{n}{2}</math> or <math>\frac{1}{2}n</math> <math>2 + 1n</math> for <math>2 + n</math> <math>2 + 0n</math> for <math>2</math></p> <p>Within a question that demands simplification, do not accept as part of a final answer involving algebra. Accept within a method when awarding partial credit, or within an explanation or general working.</p> <p>✗ Embedded values given when solving equations eg in solving <math>3x + 2 = 32</math>, <math>3 \times 10 + 2 = 32</math> for <math>x = 10</math></p> <p>To avoid penalising the two types of error below more than once within each question, do not award the mark for the <i>first</i> occurrence of each type within each question. Where a question part carries more than one mark, only the final mark should be withheld.</p>
<p>✓ Words used to precede or follow equations or expressions eg <math>t = n + 2</math> tiles or tiles = <math>t = n + 2</math> for <math>t = n + 2</math></p>	<p>! Words or units used within equations or expressions eg <math>n</math> tiles + 2 <math>n</math> cm + 2</p> <p>Do not accept on their own. Ignore if accompanying an acceptable response.</p>
<p>✓ Unambiguous letters used to indicate expressions eg <math>t = n + 2</math> for <math>n + 2</math></p>	<p>✗ Ambiguous letters used to indicate expressions eg <math>n = n + 2</math> for <math>n + 2</math></p>

<b>Responses involving coordinates</b> For example: (5, 7)	
<b>Accept ✓</b>	<b>Do not accept ✗</b>
✓ Unconventional notation eg (05, 07) (five, seven) $\begin{matrix} x & y \\ (5, & 7) \end{matrix}$ (x=5, y=7)	✗ Incorrect or ambiguous notation eg (7, 5) $\begin{matrix} y & x \\ (7, & 5) \end{matrix}$ (5x, 7y) (5 <sup>x</sup> , 7 <sup>y</sup> ) (x - 5, y - 7)

<b>Responses involving negative numbers</b> For example: -2	
<b>Accept ✓</b>	<b>Do not accept ✗</b>
	To avoid penalising the error below more than once within each question, do not award the mark for the <i>first</i> occurrence of the error within each question. Where a question part carries more than one mark, only the final mark should be withheld.  ✗ Incorrect notation eg 2-



**Responses involving probability**

A numerical probability should be expressed as a decimal, fraction or percentage only.

For example: 0.7 or  $\frac{7}{10}$  or 70%

<b>Accept ✓</b>	<b>Take care ! Do not accept ✗</b>
<p>✓ Equivalent decimals, fractions and percentages</p> <p>eg 0.700, <math>\frac{70}{100}</math>, <math>\frac{35}{50}</math>, 70.0%</p> <p>✓ A probability correctly expressed in one acceptable form which is then incorrectly converted, but is still less than 1 and greater than 0</p> <p>eg <math>\frac{70}{100} = \frac{18}{25}</math></p>	<p>The first <b>four</b> categories of error below should be ignored if accompanied by an acceptable response, but should not be accepted on their own. However, to avoid penalising the first <b>three</b> types of error below more than once within each question, do not award the mark for the <i>first</i> occurrence of each type of error unaccompanied by an acceptable response. Where a question part carries more than one mark, only the final mark should be withheld.</p> <p>! A probability that is incorrectly expressed</p> <p>eg 7 in 10 7 over 10 7 out of 10 7 from 10</p> <p>! A probability expressed as a percentage without a percentage sign</p> <p>! A fraction with other than integers in the numerator and/or denominator</p> <p>! A probability expressed as a ratio</p> <p>eg 7 : 10, 7 : 3, 7 to 10</p> <p>✗ A probability greater than 1 or less than 0</p>

### Recording marks awarded on the test paper

All questions, even those not attempted by the pupil, will be marked, with a 1 or a 0 entered in each marking space. Where 2m can be split into 1m gained and 1m lost, with no explicit order, then this will be recorded by the marker as 1  
0

The total marks awarded for a double page will be written in the box at the bottom of the right-hand page, and the total number of marks obtained on the paper will be recorded on the front of the test paper.











A total of 120 marks is available in each of tiers 3–5 and 4–6.

A total of 121 marks is available in each of tiers 5–7 and 6–8.

### Awarding levels

The sum of the marks gained on paper 1, paper 2 and the mental mathematics paper determines the level awarded. Level threshold tables, which show the mark ranges for the award of different levels, will be available on the QCA website [www.qca.org.uk/](http://www.qca.org.uk/) from Monday 20 June 2005. QCA will also send a copy to each school in July.

Schools will be notified of pupils' results by means of a marksheet, which will be returned to schools by the external marking agency with the pupils' marked scripts. The marksheet will include pupils' scores on the test papers and the levels awarded.

Tier & Question					4 by 4 grid	
3-5	4-6	5-7	6-8			
1					Correct response	Additional guidance
a				1m	<p>Correctly divides the square into quarters in a different way from the given example</p> <p>eg</p> <ul style="list-style-type: none"> <li>▪ </li> <li>▪ </li> <li>▪ </li> <li>▪ </li> </ul>	<p>! <i>Throughout the question, lines not ruled or accurate, or lines not using the intersections of the grid</i> Accept provided the pupil's intention is clear</p> <p>! <i>Throughout the question, quarters or eighths are not congruent</i> Accept provided the intention is clear for all pieces to have the same area eg, for part (a) accept</p> <ul style="list-style-type: none"> <li>♦ </li> <li>♦ </li> </ul> <p>eg, for part (b) accept</p> <ul style="list-style-type: none"> <li>♦ </li> </ul>
b				1m	<p>Correctly divides the square into eighths</p> <p>eg</p> <ul style="list-style-type: none"> <li>▪ </li> <li>▪ </li> <li>▪ </li> </ul>	

Tier & Question									<b>Heating</b>	
3-5	4-6	5-7	6-8							
<b>2</b>							<b>Correct response</b>		<b>Additional guidance</b>	
a					<b>1m</b>	Indicates the correct times in the correct order eg <ul style="list-style-type: none"> <li>■ 6 and 9:30</li> </ul>	<b>✓ Indication of morning</b> eg <ul style="list-style-type: none"> <li>♦ 6 am and 9:30 am</li> </ul> <b>! Times not accurate</b> Accept $\pm$ 5 minutes of the correct times eg, for 9:30 accept <ul style="list-style-type: none"> <li>♦ 9:25 to 9:35 inclusive</li> </ul> <b>! Use of 'half'</b> Accept colloquial use of 'half' or $\frac{1}{2}$ eg, for 9:30 accept <ul style="list-style-type: none"> <li>♦ Half (or <math>\frac{1}{2}</math>) 9</li> </ul> Do not accept an incorrect time eg, for 9:30 do not accept <ul style="list-style-type: none"> <li>♦ 9 half (or <math>\frac{1}{2}</math>)</li> </ul> <b>✗ Time(s) incorrect</b> eg <ul style="list-style-type: none"> <li>♦ 6 pm and 9:30</li> <li>♦ 6 and 21:30</li> <li>♦ 6 and 9.5</li> </ul>			
						<b>1m</b>	$3\frac{1}{2}$ or equivalent	<b>! Follow through from the first mark</b> Accept as the time interval between their two times, provided their answer is not a whole number of hours  <b>! 'Half' in words</b> Condone eg, accept <ul style="list-style-type: none"> <li>♦ 3 and a half</li> </ul>		
b					<b>2m</b>	Indicates only 17(:00) and 23(:00) correctly on the diagram, with no incorrect times shown	<b>! Positions not accurate</b> Accept provided the pupil's intention is clear			
					<i>or</i> <b>1m</b>	Indicates either 17(:00) or 23(:00) correctly on the diagram, with not more than one error  or  Indicates any two times on the diagram with a difference of 6 hours	<b>! Arrows do not indicate 'on' or 'off'</b> For 2m, condone unless the times are incorrectly labelled as 'on' or 'off' In this case, mark as 1, 0 For 1m, ignore any labels			

Tier & Question					Tickets	
3-5	4-6	5-7	6-8			
3					<b>Correct response</b>	<b>Additional guidance</b>
a				1m	5	<p><b>✗ For the first mark, £5</b></p> <p><b>! Values not rounded</b>                      Penalise only the first occurrence, even if the non-integer part is incorrect                      eg, for parts (a) and (b)</p> <ul style="list-style-type: none"> <li>• 5.2(...) or 5.3</li> <li>6.8(...) or 6.9</li> </ul> Mark as 0, 1
b				1m	6	
c				1m	£ 22	
				(U1)		

Tier & Question					Unit			
3-5	4-6	5-7	6-8					
4					<b>Correct response</b>		<b>Additional guidance</b>	
a				1m	Indicates grams	✓ <i>Unambiguous indication</i>		
				1m	Indicates litres	! <i>For both responses, correct but less suitable units indicated</i> Mark responses of kilograms then millilitres as 0, 1		
b				1m	Indicates one of the given units not credited in their (a), and gives an example of something it could measure eg <ul style="list-style-type: none"> <li>■ Use metres to measure the distance of a running track</li> <li>■ Use millimetres to measure the length of a ruler</li> <li>■ Use kilograms to measure the mass of a person [only if kilograms not given for the first mark in (a)]</li> <li>■ Use millilitres to measure the volume of drink in a can [only if millilitres not given for the second mark in (a)]</li> <li>■ Use grams to measure the mass of a piece of cheese [only if grams not given for the first mark in (a)]</li> <li>■ Use litres to measure the capacity of water in a swimming pool [only if litres not given for the second mark in (a)]</li> </ul>	! <i>Imprecise description of the property to be measured</i> Condone provided the pupil's intention is clear eg, accept <ul style="list-style-type: none"> <li>♦ Use metres to measure the size of a garden</li> <li>♦ Use millilitres to measure the amount/quantity of drink in a can</li> <li>♦ Use kilograms to measure the weight of a person</li> </ul> ! <i>Units for the correct property given, but not the most suitable for their example</i> Condone eg, accept <ul style="list-style-type: none"> <li>♦ Use millilitres to measure the volume of water in a swimming pool</li> </ul> ! <i>Property given with object unspecified or omitted</i> Condone eg, accept <ul style="list-style-type: none"> <li>♦ Use millimetres to measure the length of something</li> <li>♦ Use kilograms to measure the mass</li> </ul> ✗ <i>Object given without explicit indication of the property to be measured</i> eg <ul style="list-style-type: none"> <li>♦ Use millimetres to measure a ruler</li> <li>♦ Use kilograms to measure a person</li> </ul> ✗ <i>Units used that are not from the given list</i> eg <ul style="list-style-type: none"> <li>♦ Use centimetres to measure the length of a ruler</li> </ul>		

U1





Tier & Question					Half price	
3-5	4-6	5-7	6-8			
<b>6</b>					<b>Correct response</b>	<b>Additional guidance</b>
a				1m	£ 2.84	
b				1m	£ 13.98	

Tier & Question					Teachers	
3-5	4-6	5-7	6-8			
<b>7</b>					<b>Correct response</b>	<b>Additional guidance</b>
a				1m	187 860	
b				1m	1350	<b>x</b> -1350

Tier & Question					Membership	
3-5	4-6	5-7	6-8			
<b>8</b>	<b>1</b>				<b>Correct response</b>	<b>Additional guidance</b>
a	a			1m	October	<p>✓ <i>Unambiguous indication of month</i> eg • O</p> <p>! <i>Correct frequency of 32 given</i> Ignore alongside indication of the correct month, but do not accept on its own</p>
b	b			1m	11	

Tier & Question					Factor	
3-5	4-6	5-7	6-8			
9	2				Correct response	Additional guidance
a	a			1m	<p>Indicates Yes and gives a correct explanation</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ <math>3 \times 10 = 30</math></li> <li>■ <math>30 \div 3 = 10</math></li> <li>■ 30 is a multiple of 3</li> <li>■ 3 goes into 30 exactly</li> <li>■ 30 is in the 3 times table</li> </ul>	<p>✓ <i>Minimally acceptable explanation</i></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ <math>3 \times 10</math></li> <li>♦ <math>30 \div 3</math> has no remainder</li> <li>♦ 30 divides by 3</li> <li>♦ 3 goes into 30</li> <li>♦ <math>30 \div 10</math></li> <li>♦ <math>3 + 0 = 3</math> which is in the 3 times table</li> </ul> <p>! <i>Use of repeated addition</i></p> <p>Condone</p> <p>eg, accept</p> <ul style="list-style-type: none"> <li>♦ Keep going up in 3s and you get to 30</li> </ul> <p>! <i>Use of 'it' or other ambiguous language</i></p> <p>Condone provided either 3 or 30 is used, implying 'it' is the other number</p> <p>eg, accept</p> <ul style="list-style-type: none"> <li>♦ 30 divides by it</li> <li>♦ The lower number goes into it</li> <li>♦ It's in the 3 times table</li> </ul> <p>eg, do not accept</p> <ul style="list-style-type: none"> <li>♦ It goes into it 10 times</li> </ul> <p>! <i>Response contains an incorrect statement</i></p> <p>Ignore alongside a correct response</p> <p>eg, accept</p> <ul style="list-style-type: none"> <li>♦ 30 divides by 3 as 3 is a multiple of 30</li> </ul> <p>eg, do not accept</p> <ul style="list-style-type: none"> <li>♦ <math>3 \div 30 = 10</math></li> <li>♦ 30 goes into 3 exactly</li> </ul> <p>✗ <i>Incomplete or incorrect explanation</i></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ 3 is a factor of 30</li> <li>♦ <math>30 \div 3</math></li> <li>♦ It adds up to 30</li> <li>♦ They're both in the 3 times table</li> <li>♦ Because there is a 3 in it</li> </ul>
b	b			1m	<p>Gives a factor of 30 greater than 3, ie</p> <p style="text-align: center;">5, 6, 10, 15 or 30</p>	

U1

Tier & Question						<b>Shapes on a grid</b>	
3-5	4-6	5-7	6-8				
10	3				Correct response	Additional guidance	
a	a			1m	20		
b	b			1m	60	<p><b>!</b> <i>Follow through</i> Accept follow through as their (a) × 3, provided their (a) was not 5</p>	
c	c			1m	4	<p><b>!</b> <i>Operation repeated</i> eg ♦ × 4 Condone</p> <p><b>×</b> <i>More than one number given</i> eg ♦ 2 × 2</p>	

U1





Tier & Question						<b>Meal</b>	
3-5	4-6	5-7	6-8				
11	4				Correct response	Additional guidance	
				2m	£ 276		
				or 1m	Shows the digits 276 eg ■ 2.76		
					or Shows the value 23, with no evidence of an incorrect method	<p><b>×</b> <i>For 1m, incorrect method</i> eg ♦ 11 + 12 = 23</p>	
					or Shows or implies a complete correct method with not more than one computational or rounding error eg ■ $\frac{253}{11} \times 12$ ■ $253 \div 11 = 13$ (error) $253 + 13 = 266$ ■ $12 \div 11 = 1.09(\dots)$ , $1.09$ (premature rounding) $\times 253 = 275.77$		

Tier & Question										<b>Rhombus area</b>																															
3-5	4-6	5-7	6-8																																						
12	5					Correct response		Additional guidance																																	
a	a			1m	10.2 to 10.4 inclusive and 6.6 to 6.8 inclusive, in either order	<p>✓ <i>Throughout the question, equivalent fractions or decimals</i></p> <p>✓ <i>Follow through as the product of their two values for part (a) ÷ 2</i> As this is an algebra mark, accept follow through from whole numbers as well as decimals</p> <p>! <i>For part (b), their value rounded</i> Accept correct rounding to the nearest integer or better, or truncation to one decimal place or better Do not accept incorrect rounding or truncation to an integer unless a correct method or a more accurate value is seen</p> <p>Markers may find the following values for the diagonals and corresponding areas useful:</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th></th> <th colspan="4" style="text-align: center;">(error)</th> </tr> <tr> <th></th> <th>6.5</th> <th>6.6</th> <th>6.7</th> <th>6.8</th> </tr> </thead> <tbody> <tr> <td>10.2</td> <td>33.15</td> <td>33.66</td> <td>34.17</td> <td>34.68</td> </tr> <tr> <td>10.3</td> <td>33.475</td> <td>33.99</td> <td>34.505</td> <td>35.02</td> </tr> <tr> <td>10.4</td> <td>33.8</td> <td>34.32</td> <td>34.84</td> <td>35.36</td> </tr> <tr> <td>10.5</td> <td>34.125</td> <td>34.65</td> <td>35.175</td> <td>35.7</td> </tr> </tbody> </table> <p>(error)</p>							(error)					6.5	6.6	6.7	6.8	10.2	33.15	33.66	34.17	34.68	10.3	33.475	33.99	34.505	35.02	10.4	33.8	34.32	34.84	35.36	10.5	34.125	34.65	35.175	35.7
	(error)																																								
	6.5	6.6	6.7	6.8																																					
10.2	33.15	33.66	34.17	34.68																																					
10.3	33.475	33.99	34.505	35.02																																					
10.4	33.8	34.32	34.84	35.36																																					
10.5	34.125	34.65	35.175	35.7																																					
b	b			1m	<p>Gives the correct area using their values for the lengths of the diagonals in part (a)</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ From 10.3 and 6.7 in part (a), area of 34.505 (or 3450.5)</li> </ul> <p>or</p> <p>Gives the correct area using two values seen in part (b), even if they are different from their values for the lengths of the diagonals in part (a)</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ From 10 and 7 seen in part (b), area of 35</li> </ul>																																				
				1m	<p>Shows the correct unit for their area</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ 34.505 cm<sup>2</sup></li> <li>■ 3450.5 mm<sup>2</sup></li> <li>■ Product of their two values for part (a) ÷ 2 and cm<sup>2</sup> seen</li> <li>■ Product of their two values for part (a) ÷ 2 × 100 and mm<sup>2</sup> seen</li> </ul>	<p>! <i>Area not followed through from their (a) or omitted, but units given</i> If the first mark in part (b) for their correct area has not been awarded, condone either cm<sup>2</sup> or mm<sup>2</sup> seen for the second mark in part (b)</p>																																			

Tier & Question									<b>Mobile phones</b>	
3-5	4-6	5-7	6-8							
13	6						<b>Correct response</b>			<b>Additional guidance</b>
					1m		Gives a value between 1 and 2 inclusive			<b>! 'Million' repeated</b> eg, for the first mark <ul style="list-style-type: none"> <li>♦ <math>1\frac{1}{2}</math> million</li> <li>♦ 1 500 000</li> </ul> Condone
					1m		Gives a value between 49.5 and 50.5 inclusive			
					1m		Gives a value between 10 and 12 inclusive			




Tier & Question									<b>Arranging numbers</b>	
3-5	4-6	5-7	6-8							
14	7						<b>Correct response</b>			<b>Additional guidance</b>
					2m		Gives both correct ways that are different from the example given, ie  <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; border-radius: 10px; padding: 5px; text-align: center;">2 , 3</div> <div style="border: 1px solid black; border-radius: 10px; padding: 5px; text-align: center;">1 , 4 , 5</div> </div> <p style="text-align: center;">and</p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; border-radius: 10px; padding: 5px; text-align: center;">1 , 4</div> <div style="border: 1px solid black; border-radius: 10px; padding: 5px; text-align: center;">2 , 3 , 5</div> </div>			<b>! Operations given</b> Ignore eg, for 2, 3 accept <ul style="list-style-type: none"> <li>♦ 2 + 3</li> </ul> <b>! First and second groups transposed within an otherwise completely correct response [answer lines ignored]</b> eg <ul style="list-style-type: none"> <li>♦</li> </ul> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; border-radius: 10px; padding: 5px; text-align: center;">1, 4, 5</div> <div style="border: 1px solid black; border-radius: 10px; padding: 5px; text-align: center;">2, 3</div> </div> <p style="text-align: center;">and</p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; border-radius: 10px; padding: 5px; text-align: center;">2, 3, 5</div> <div style="border: 1px solid black; border-radius: 10px; padding: 5px; text-align: center;">1, 4</div> </div> Mark as 0, 1
					or 1m		Gives one of the two correct ways that are different from the example given			

U1

Tier & Question					What shape?	
3-5	4-6	5-7	6-8			
15	8				<b>Correct response</b>	<b>Additional guidance</b>
a	a			1m	Draws a triangle with no right angle eg ■ 	! <i>Lines not ruled or accurate</i> Accept provided the pupil's intention is clear  ! <i>Vertices not on grid intersections</i> Accept provided it is clear that the conditions have been satisfied
b	b			1m	Draws a quadrilateral with no right angles eg ■  ■  ■ 	
c	c			1m	Indicates 1	✓ <i>Unambiguous indication including angle marked on diagram</i>

Tier & Question				Refer to the new algebra general guidance	Algebra grids
3-5	4-6	5-7	6-8		
17	9	1		Correct response	Additional guidance
			1m	<p>Completes the grid correctly, giving simplified expressions, ie</p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; border-radius: 10px; padding: 5px 15px;"><math>8k</math></div> <div style="border: 1px solid black; border-radius: 10px; padding: 5px 15px;"><math>3k</math></div> </div> <div style="display: flex; justify-content: center; align-items: center; margin-top: 10px;"> <div style="border: 1px solid black; border-radius: 10px; padding: 5px 15px;"><math>11k</math></div> </div>	
			2m	<p>Completes the grid correctly, giving simplified expressions eg</p> <ul style="list-style-type: none"> <li>■</li> </ul> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="border: 1px solid black; border-radius: 10px; width: 40px; height: 20px; margin-right: 10px;"></div> <div style="border: 1px solid black; border-radius: 10px; width: 40px; height: 20px; margin-right: 10px;"></div> <div style="border: 1px solid black; border-radius: 10px; padding: 5px 15px;"><math>3a + 3b</math></div> </div> <div style="display: flex; justify-content: space-around; align-items: center; margin-top: 10px;"> <div style="border: 1px solid black; border-radius: 10px; padding: 5px 15px;"><math>6a + 5b</math></div> <div style="border: 1px solid black; border-radius: 10px; width: 40px; height: 20px;"></div> </div> <div style="display: flex; justify-content: center; align-items: center; margin-top: 10px;"> <div style="border: 1px solid black; border-radius: 10px; padding: 5px 15px;"><math>13a + 10b</math></div> </div>	
			or 1m	Gives two correct simplified expressions	<p><b>! For 1m, follow through</b> Accept follow through from their incorrect expression for <math>6a + 5b</math>, provided their incorrect expression contains only a term in <math>a</math> and a term in <math>b</math></p>



Tier & Question						<b>1976 v 2002</b>	
3-5	4-6	5-7	6-8				
16	10	2		Correct response		Additional guidance	
a	a	a		1m	£ 4		
b	b	b		2m	<p>Completes the pie chart correctly eg</p> 	<p><b>! Labels abbreviated</b> Accept unambiguous indications of category names eg, for 2m accept</p> 	
				or 1m	<p>Draws all four sectors correctly but fails to label or labels incorrectly</p> <p>or</p> <p>Draws and labels any two of the sectors correctly</p> <p>or</p> <p>Makes an error in drawing either the rent or the food sector provided rent sector &gt; food sector, and follows through correctly to divide the remaining space into two equal sectors for entertainment and other</p>	<p>Do not accept amounts of money as the only labels, but ignore alongside correct labels</p> <p><b>! Lines not ruled or accurate</b> Accept provided the pupil's intention is clear</p> <p><b>✗ Sector not continuous</b> Do not accept as a correct sector eg, for the rent sector do not accept</p> 	

Tier & Question						Correct response	Additional guidance																
3-5	4-6	5-7	6-8																				
18	11	3																					
					2m	<p>Indicates the village shop and gives a correct justification, based on correctly calculating a pair of comparable values eg</p> <ul style="list-style-type: none"> <li>■ At the supermarket <math>6.25 \times 6 = 37.5(0)</math> At the village shop <math>7.20 \times 5 = 36</math></li> <li>■ <math>6.25 \times 6 - 7.2 \times 5 = 1.5</math></li> <li>■ <math>6.25 \div 5 = 1.25</math>, <math>7.20 \div 6 = 1.2(0)</math></li> <li>■ £75 for 60 or £72 for 60</li> <li>■ For £1 you get <math>\frac{4}{5}</math> of a pen or <math>\frac{5}{6}</math> of a pen</li> <li>■ You pay 95p extra for 1 more pen, but they're at least £1.20 each so it must be a better deal</li> </ul>	<p><b>✗ For 2m, no decision</b></p> <p><b>✓ For 2m, correct decision and any pair of comparable values shown</b> Note that common pairs (in pounds) are:</p> <table style="width: 100%; border: none;"> <tr> <td>37.5 and 36</td> <td style="text-align: right;">(per 30 pens)</td> </tr> <tr> <td>1.25 and 1.2</td> <td style="text-align: right;">(per 1 pen)</td> </tr> <tr> <td>6.25 and 6</td> <td style="text-align: right;">(per 5 pens)</td> </tr> <tr> <td>7.5 and 7.2</td> <td style="text-align: right;">(per 6 pens)</td> </tr> <tr> <td>75 and 72</td> <td style="text-align: right;">(per 60 pens)</td> </tr> <tr> <td>18.75 and 18</td> <td style="text-align: right;">(per 15 pens)</td> </tr> <tr> <td>0.95 and 1.2 [or 1.25]</td> <td style="text-align: right;">(1 extra pen)</td> </tr> <tr> <td>0.8 and 0.83(...)</td> <td style="text-align: right;">(pens per pound)</td> </tr> </table> <p><b>! For 2m or 1m, comparison is per 5 pens or per 6 pens but the given price is not restated</b> Condone eg, for 2m accept</p> <ul style="list-style-type: none"> <li>• At the supermarket, 6 pens would be £7.50</li> </ul> <p><b>! Additional incorrect working</b> Ignore</p>	37.5 and 36	(per 30 pens)	1.25 and 1.2	(per 1 pen)	6.25 and 6	(per 5 pens)	7.5 and 7.2	(per 6 pens)	75 and 72	(per 60 pens)	18.75 and 18	(per 15 pens)	0.95 and 1.2 [or 1.25]	(1 extra pen)	0.8 and 0.83(...)	(pens per pound)
37.5 and 36	(per 30 pens)																						
1.25 and 1.2	(per 1 pen)																						
6.25 and 6	(per 5 pens)																						
7.5 and 7.2	(per 6 pens)																						
75 and 72	(per 60 pens)																						
18.75 and 18	(per 15 pens)																						
0.95 and 1.2 [or 1.25]	(1 extra pen)																						
0.8 and 0.83(...)	(pens per pound)																						
					or 1m	<p>Shows a correct pair of comparable values but makes either an incorrect or no decision</p> <p>or</p> <p>Shows a complete correct method for finding a pair of comparable values with not more than one computational or rounding error, and follows through to make their correct decision eg</p> <ul style="list-style-type: none"> <li>■ <math>6 \times 6.25</math>, <math>5 \times 7.20</math> [village shop indicated]</li> <li>■ <math>6.25 \div 5 = 1.05</math> (error), <math>7.20 \div 6 = 1.20</math> [supermarket indicated]</li> </ul> <p>or</p> <p>Makes a correct decision but the justification uses only the difference between a pair of comparable values eg</p> <ul style="list-style-type: none"> <li>■ The packs of 6 would be £1.50 cheaper</li> <li>■ A pen is 5p cheaper</li> </ul>																	

U1

Tier & Question						<b>Counters</b>	
3-5	4-6	5-7	6-8				
20	12	4		Correct response		Additional guidance	
a	a	a		1m	$\frac{1}{3}$ or equivalent probability	! <i>Value rounded</i> Accept 0.33 or better, or the percentage equivalents	
b	b	b		1m	3		

Tier & Question				<i>Marking overlay available</i>		<b>From London</b>	
3-5	4-6	5-7	6-8				
19	13	5		Correct response		Additional guidance	
a	a	a		1m	$160 \pm 2$		
b	b	b		1m	$350 \pm 5$		
c	c	c		2m	Indicates the correct position of Madrid within the tolerance as shown on the overlay	! <i>For 2m, Madrid not labelled</i> Condone provided the intended position is clear	
				or 1m	Indicates an angle of $195^\circ \pm 2^\circ$ clockwise from north, within the tolerance as shown on the overlay	! <i>For 1m, angle indicated with a short line</i> Accept provided the angle is within the tolerance as shown on the overlay, were the line to be extended	
					or Shows a length of $6.5\text{cm} \pm 0.2\text{cm}$ , within the tolerance as shown on the overlay, even if it is incorrectly positioned	! <i>For 1m, angle or length indicated by a point without a line joined to London</i> Accept provided the angle or length is within the tolerance as shown on the overlay	


Tier & Question						<b>How many?</b>					
3-5	4-6	5-7	6-8								
21	14	6		Correct response		Additional guidance					
a	a	a		1m	<p>Gives the correct number of boys and girls, ie</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <th>Number of boys</th> <th>Number of girls</th> </tr> <tr> <td style="text-align: center;">.....18.....</td> <td style="text-align: center;">.....9.....</td> </tr> </table>	Number of boys	Number of girls	.....18.....	.....9.....	<p><b>!</b> <i>Numbers correct but numbers of boys and girls transposed</i>                      Penalise only the first occurrence                      eg, for all three parts</p> <ul style="list-style-type: none"> <li>• 9, 18</li> <li>13, 15</li> <li>18, 9</li> </ul> <p>Mark as 0, 1, 1</p> <p><b>!</b> <i>Values given as tallies</i>                      Condone provided they are grouped in fives</p>	
Number of boys	Number of girls										
.....18.....	.....9.....										
b	b	b		1m	<p>Gives the correct number of boys and girls, ie</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <th>Number of boys</th> <th>Number of girls</th> </tr> <tr> <td style="text-align: center;">.....15.....</td> <td style="text-align: center;">.....13.....</td> </tr> </table>	Number of boys	Number of girls	.....15.....	.....13.....		
Number of boys	Number of girls										
.....15.....	.....13.....										
c	c	c		1m	<p>Gives the correct number of boys and girls, ie</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <th>Number of boys</th> <th>Number of girls</th> </tr> <tr> <td style="text-align: center;">.....9.....</td> <td style="text-align: center;">.....18.....</td> </tr> </table>	Number of boys	Number of girls	.....9.....	.....18.....		
Number of boys	Number of girls										
.....9.....	.....18.....										



Tier & Question								<b>Tennis prizes</b>	
3-5	4-6	5-7	6-8						
				<b>Correct response</b>		<b>Additional guidance</b>			
17	9	1		2m	<p>Indicates France and gives a correct justification</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ <math>1000\ 000 \div 2.7 = 370\ 370.(...)</math>, <math>780\ 000 \div 1.54 = 506\ 493.(...)</math></li> <li>■ <math>\frac{1000\ 000}{2.7} &lt; \frac{780\ 000}{1.54}</math></li> <li>■ <math>1000\ 000 \div 2.7 \times 1.54 = 570\ 370.(...)</math></li> <li>■ <math>780\ 000 \div 1.54 \times 2.7 = 1\ 367\ 532.(...)</math></li> </ul>	<p>✓ <i>For 2m, minimally acceptable justification</i></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ 370 370 and 506 493 (or 506 494) seen</li> <li>♦ <math>\frac{1000\ 000}{2.7}</math>, <math>\frac{780\ 000}{1.54}</math></li> <li>♦ <math>1000\ 000 \div 270 = 3703.(...)</math> (or 3704), <math>780\ 000 \div 154 = 5064.(...)</math> (or 5065)</li> <li>♦ 570 370.(...) seen</li> <li>♦ 1 367 532.(...) seen</li> </ul> <p>! <i>Values rounded or estimated</i></p> <p>For 2m, accept values of 370 0(00) and 500 0(00) or better, 570 000 or better, or 1 400 000 or better</p> <p>Accept other estimates only if a correct method or a more accurate value is seen</p> <p>eg, accept</p> <ul style="list-style-type: none"> <li>♦ £1 is about <math>2\frac{1}{2}</math> dollars, so 1000 000 dollars is about £400 000, £1 is about <math>1\frac{1}{2}</math> euros, so 780 000 euros is about £500 000</li> </ul> <p>✗ <i>For 2m or 1m, justification simply repeats the decision made</i></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ 1000 000 Australian dollars are less than 780 000 euros</li> </ul>			
				or 1m	<p>Indicates France and gives a partial justification</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ <math>1000\ 000 \approx \pounds 400\ 000</math>, <math>780\ 000 \approx \pounds 500\ 000</math></li> <li>■ Australia: 370 France: 506 [values truncated with no indication of method or that original values were of the same magnitude]</li> </ul> <p>or</p> <p>Gives a correct justification but makes an incorrect or no decision</p> <p>or</p> <p>Gives a correct justification with not more than one computational or rounding error, but follows through to make their correct decision</p>				
				(U1)					

Tier & Question				<i>Marking overlay available</i>		<b>Enlargement</b>
3-5	4-6	5-7	6-8			
	<b>18</b>	<b>10</b>	<b>2</b>		<b>Correct response</b>	<b>Additional guidance</b>
				2m	Draws the correct enlargement with vertices within the tolerances as shown on the overlay	<p><b>! <i>Lines not ruled or accurate</i></b> Accept provided the pupil's intention is clear</p> <p><b>! <i>Construction lines shown</i></b> Ignore</p> <p><b>✗ <i>Enlargement is the correct size but in an incorrect orientation</i></b></p>
				or 1m	<p>Within an otherwise correct enlargement, the only error is that the vertices are not correctly joined</p> <p>or</p> <p>Their enlargement is the correct size and orientation as shown by the overlay, with vertices joined correctly, but is in the incorrect position</p>	



Tier & Question								<b>Heron of Alexandria</b>	
3-5	4-6	5-7	6-8						
	<b>19</b>	<b>11</b>	<b>3</b>			<b>Correct response</b>		<b>Additional guidance</b>	
				2m	$\sqrt{56}$ , $2\sqrt{14}$ , 7.48(...) or 7.5, with no evidence of an incorrect method			✓ <i>Equivalent fractions or decimals</i>  ! <i>For 2m, answer of 7</i> Do not accept unless a correct method or a more accurate value is seen  ✗ <i>Incorrect method</i> eg ♦ $3 \times 5 \div 2 = 7.5$ ♦	
				or 1m	Shows or implies at least two of the following three correct steps 1. Shows or implies that the value of $s$ is 7 2. Substitutes correctly the values of $a$ , $b$ and $c$ and their $s$ into the expression $s(s - a)(s - b)(s - c)$ 3. Takes the square root of the correct result of their substitution eg ■ 56 seen [step 3 omitted] ■ $7(7 - 3)(7 - 5)(7 - 6)$ [step 3 omitted] ■ $\sqrt{7 \times 4 \times 2 \times 2}$ ( <i>error</i> ) = 10.5(...) or 10.6 [step 2 incorrect] ■ $\sqrt{14(14 - 3)(14 - 5)(14 - 6)}$ = 105.(...) [step 1 incorrect] ■ 7.4 [correct value truncated]				
					or  Shows the value 51, 51.3(...) or 51.4 [the only error is to use $s$ as 11]				
					or  Shows the value 21, 21.1(...) or 21.2 [the only error is to take the square root of 7 before multiplying by 4 and 2]				

Tier & Question						<b>Hands</b>	
3-5	4-6	5-7	6-8				
20	12	4			Correct response		Additional guidance
a	a	a	1m		$\frac{7}{15}$ or equivalent probability		<p><b>! Value rounded or truncated</b> Accept 0.46(...) or 0.47 or the percentage equivalents Do not accept 0.5 unless a correct method or a more accurate value is seen</p>
b	b	b	1m		$\frac{1}{10}$ or equivalent probability		<p><b>! Follow through</b> Accept follow through from an incorrect total number of pupils seen in part (a), provided their total is not 4, 16 or 27 eg, from <math>\frac{14}{29}</math> for part (a) accept</p> <ul style="list-style-type: none"> <li>• <math>\frac{3}{29}</math></li> </ul>
c	c	c	1m		$\frac{2}{3}$ or equivalent probability		<p><b>! Value rounded</b> Accept 0.66(...) or 0.67 or the percentage equivalents</p>

Tier & Question						<b>Screens</b>	
3-5	4-6	5-7	6-8				
21	13	5			Correct response		Additional guidance
			1m		8		<p><b>! Values transposed but otherwise correct</b> Mark as 0, 1</p> <p><b>! The only error is to work with ratios that are prematurely rounded</b> For the first value between 7.65 and 8.1 inclusive (excluding 8), and for the second value between 5.85 and 6.3 inclusive (excluding 6), mark as 0, 1</p>
			1m		6		

Tier & Question								<b>Spinning</b>	
3-5	4-6	5-7	6-8						
	22	14	6			<b>Correct response</b>			<b>Additional guidance</b>
				2m	0.15 or equivalent probability				<p><b>✗ For 2m, incorrect notation</b></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ <math>0.1 \frac{1}{2}</math></li> <li>♦ 0.1.5</li> </ul>
				or 1m	Shows or implies the intention to add the given probabilities, subtract the sum from 1 and then divide by 2, even if there are errors eg <ul style="list-style-type: none"> <li>■ <math>0.1 + 0.6 = 0.7</math></li> <li>■ <math>\frac{1 - 0.7}{2}</math></li> <li>■ <math>0.3 \div 2</math></li> <li>■ <math>\frac{1.5}{10}</math></li> </ul>				

Tier & Question								<b>Number</b>	
3-5	4-6	5-7	6-8						
	23	15	7			<b>Correct response</b>			<b>Additional guidance</b>
				2m	11				
				or 1m	Forms or implies a correct equation eg <ul style="list-style-type: none"> <li>■ <math>8x - 66 = 2x</math></li> <li>■ <math>6y = 66</math></li> <li>■ <math>66 \div 6</math></li> </ul>				<p><b>! Method used is trial and improvement</b> Note that no partial credit can be given</p> <p><b>! Equation involving words</b> Accept provided the operation involved in 'twice the number I was thinking of' has been interpreted eg, for 1m accept           <ul style="list-style-type: none"> <li>♦ Number <math>\times</math> 8 minus 66 = number <math>\times</math> 2</li> <li>♦ 66 is the same as 6 times the number</li> </ul>           eg, for 1m do not accept           <ul style="list-style-type: none"> <li>♦ <math>8x - 66 = \text{twice } x</math></li> </ul> </p>

U1

Tier & Question				A level results	
3-5	4-6	5-7	6-8		
24	16	8		Correct response	Additional guidance
			2m	6300	! <i>Incorrect use of % sign</i> Ignore
			<i>or</i> 1m	Shows the digits 63(00)  or  Shows the value 13 680 or 19 980  or  Shows the digits 1368(0) and 1998(0)  or  Shows a complete correct method with not more than one computational error eg <ul style="list-style-type: none"> <li>■ <math>\frac{37}{100} \times 54\,000 - \frac{19}{100} \times 72\,000</math></li> <li>■ <math>37 \times 540 - 19 \times 720</math></li> </ul>	

Tier & Question				Refer to the new algebra general guidance	Solutions
3-5	4-6	5-7	6-8		
	25	17	9		
		a	a	1m	
					<p>Indicates No and gives a correct explanation</p> <p>The most common correct explanations:</p> <p>Show that the two sides of the equation are not equal when <math>y = 17</math></p> <p>eg</p> <ul style="list-style-type: none"> <li>■ <math>14 \times 17 - 51 = 187</math>, but <math>187 + 4 \times 17 = 255</math></li> <li>■ <math>14y - 51 = 187</math>, so it will go over when you add the <math>4y</math></li> <li>■ The equation simplifies to <math>10y = 238</math>, but <math>10 \times 17 = 170</math></li> </ul> <p>Show the correct solution or show a correct method for solving the equation that demonstrates that the solution cannot be 17</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ <math>14y - 51 = 187 + 4y</math>  <math>10y = 238</math>  <math>y = 23.8</math></li> <li>■ <math>(187 + 51) \div 10 \neq 17</math></li> </ul> <p>Show or imply that <math>y = 17</math> is a correct solution to <math>14y - 51 = 187</math></p> <p>eg</p> <ul style="list-style-type: none"> <li>■ <math>14 \times 17 - 51 = 187</math>, but there is another <math>4 \times 17</math> to add to the 187 on the other side</li> </ul>
					<p><b>✓ Minimally acceptable explanation</b></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ <math>187 \neq 255</math></li> <li>♦ <math>14 \times 17 - 51 \neq 187 + 4 \times 17</math></li> <li>♦ <math>14 \times 17 - 51 = 187</math> so you don't need <math>4y</math></li> <li>♦ <math>14y - 51 = 187 + 0</math></li> </ul> <p><b>✗ Incomplete or incorrect explanation</b></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ When you substitute <math>y = 17</math> into both sides, you get different answers</li> <li>♦ <math>14 \times 17 - 51 = 187</math></li> <li>♦ <math>14 \times 17 - 51 = 187</math>, but <math>187 + 4 \times 17 = 225</math> (error)</li> </ul> <p><b>✓ Minimally acceptable explanation</b></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ 23.8 or equivalent seen</li> <li>♦ <math>10y = 238</math>, so <math>y \neq 17</math></li> </ul> <p><b>✗ Incorrect explanation</b></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ <math>18y = 238</math>  <math>y = 13.2</math></li> <li>♦ <math>10y = 136</math>  <math>y = 13.6</math></li> </ul> <p><b>✓ Minimally acceptable explanation</b></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ If <math>y = 17</math>, <math>14y - 51 = 187</math>, without <math>+ 4y</math></li> <li>♦ The left-hand side is 187, but the other side is 187 plus something</li> </ul> <p><b>✗ Incomplete explanation</b></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ If <math>y = 17</math>, <math>14y - 51 = 187</math></li> </ul>

Tier & Question				Refer to the new algebra general guidance	Solutions (cont)	
3-5	4-6	5-7	6-8			
	25	17	9			
		b	b	1m	<p>Indicates No and gives a correct explanation</p> <p>The most common correct explanations:</p> <p>Show that the two sides of the equation cannot be equal when <math>y = 17</math></p> <p>eg</p> <ul style="list-style-type: none"> <li>■ <math>3 \times 17^2 = 867</math>, not 2601</li> <li>■ <math>y^2 = \frac{2601}{3}</math> = 867, but <math>17 \times 17 = 289</math></li> <li>■ If <math>y = 20</math>, <math>3y^2 = 1200</math> which is still smaller than 2601, so <math>y</math> can't be 17</li> <li>■ <math>17^2</math> ends in a 9, then this number <math>\times 3</math> ends in a 7, so it can't be 2601</li> </ul> <p>Show the correct solution or show a correct method for solving the equation that demonstrates that the solution cannot be 17</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ <math>3y^2 = 2601</math> <math>y^2 = 867</math> <math>y = \pm 29.(\dots)</math></li> </ul> <p>Address the misconception</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ <math>(3 \times 17)^2 = 2601</math>, so <math>3 \times 17^2 \neq 2601</math></li> <li>■ Square 17 first, then <math>\times 3</math> and your answer is much smaller than 2601</li> </ul>	<p><b>✓ Minimally acceptable explanation</b></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ 867</li> <li>♦ <math>3 \times 289 \neq 2601</math></li> <li>♦ <math>y^2 = 867</math>, but <math>17^2 \neq 867</math></li> <li>♦ <math>17^2</math> ends in 9, then <math>\times 3</math> ends in 7</li> </ul> <p><b>✗ Incomplete explanation</b></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ <math>3 \times 17^2 \neq 2601</math></li> <li>♦ When you substitute <math>y = 17</math> into the equation, you don't get 2601</li> <li>♦ <math>3 \times 17 \times 17</math> is far too small to be 2601</li> </ul> <p><b>✓ Minimally acceptable explanation</b></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ It's <math>\pm 29.(\dots)</math></li> <li>♦ <math>\sqrt{\frac{2601}{3}} \neq 17</math></li> </ul> <p><b>! Only positive solution shown</b></p> <p>Condone</p> <p>eg, accept as minimal</p> <ul style="list-style-type: none"> <li>♦ It's 29.(\dots)</li> </ul> <p><b>✗ Incorrect explanation</b></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ <math>y^2 = 1300.5</math> <math>y = 36.(\dots)</math></li> </ul> <p><b>✓ Minimally acceptable explanation</b></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ <math>(3 \times 17)^2 = 2601</math></li> <li>♦ <math>17^2</math> then <math>\times 3 \neq 2601</math></li> <li>♦ They've squared <math>3y</math>, not just <math>y</math></li> <li>♦ You do the power, then multiply</li> <li>♦ True for <math>(3y)^2</math></li> <li>♦ <math>9y^2 = 2601</math></li> </ul> <p><b>✗ Incomplete explanation</b></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ <math>3 \times 17^2 \neq 2601</math></li> </ul>

Tier & Question				Refer to the new algebra general guidance		Simplify
3-5	4-6	5-7	6-8			
	26	18	10	1m	$9 + 2k$	
				1m	$k(k + 6)$ or $k^2 + 6k$	
				1m	$6k^2$	
				1m	$3k$	

Tier & Question								<b>Watching</b>		
3-5	4-6	5-7	6-8							
		19	12			<b>Correct response</b>			<b>Additional guidance</b>	
				2m		5 hours 12 minutes				
				<i>or</i> 1m		Shows or implies a correct method for finding the time interval for Friday, Saturday or Sunday eg <ul style="list-style-type: none"> <li>■ <math>26 \div 5</math></li> <li>■ 5.2</li> <li>■ 5 hours 20 (<i>error</i>) minutes</li> <li>■ 5 hours 2 (<i>error</i>) minutes</li> <li>■ <math>1560 \div 10 \times 2</math></li> <li>■ 312</li> </ul> or Shows or implies a correct method for finding the time interval for Monday, Tuesday, Wednesday or Thursday eg <ul style="list-style-type: none"> <li>■ 2 hours 36 minutes</li> <li>■ <math>26 \div 10</math></li> <li>■ 2.6</li> <li>■ 156</li> </ul> or Shows a correct conversion of a number of hours or minutes to hours and minutes eg <ul style="list-style-type: none"> <li>■ 1.3 hrs (<i>error</i>) = 1 hour 18 minutes</li> <li>■ 3.71(...) hrs (<i>error</i>) = 3 hours 42(...) or 43 minutes</li> <li>■ <math>1460</math> (<i>error</i>) <math>\div 5 = 292</math>, 292 mins = 4 hours 52 minutes</li> </ul>				<b>✗ For 1m, number of hours or minutes is equivalent to a multiple of <math>\frac{1}{4}</math> hour</b>
				(U1)						







Tier & Question				Sequences	
3-5	4-6	5-7	6-8	Refer to the new algebra general guidance	
		21	13	Correct response	Additional guidance
		a	a	1m 28	
		b	b	2m Gives all three correct terms in any order eg ▪ $-1, 0, \frac{1}{9}$	<p><b>!</b> <i>First two terms shown as fractions</i> eg, for the first term ♦ <math>\frac{-1}{1}</math> eg, for the second term ♦ <math>\frac{0}{4}</math> For 2m, accept provided there is no further incorrect processing</p> <p><b>!</b> <i>For 2m or 1m, <math>\frac{1}{9}</math> rounded</i> Accept 0.11 or better Do not accept 0.1 unless a correct method or a more accurate value is seen</p>
				or 1m Gives any two correct terms or Shows or implies correct substitution and interpretation of the 'squared' for all three terms, even if there is further incorrect processing eg ▪ $\frac{1-2}{1 \times 1}, \frac{2-2}{2 \times 2}, \frac{3-2}{3 \times 3}$ ▪ $-\frac{1}{1} = 1$ (error) $\frac{0}{4} = 4$ (error) $\frac{1}{9} = 0.9$ (error)	

Tier & Question				Bracket multiplication	
3-5	4-6	5-7	6-8	Correct response	Additional guidance
		22	14		
				1m Gives a correct expression without brackets eg ▪ $y^2 - 6y$	<p><b>!</b> <i>Unconventional notation</i> Condone eg, for the first mark accept ♦ <math>y \times y - y6</math></p> <p><b>✗</b> <i>Incorrect further working</i> eg, for the first mark ♦ <math>y^2 - 6y = -5y^2</math></p>
				1m Gives a correct expression without brackets eg ▪ $k^2 + 5k + 6$ ▪ $k^2 + 2k + 3k + 6$	

Tier & Question										<b>Parallelogram</b>	
3-5	4-6	5-7	6-8								
		23	15								
				1m	<p>Gives <math>h = 80</math> and gives a correct reason eg</p> <ul style="list-style-type: none"> <li>■ <math>h</math> is an alternate angle with the <math>80^\circ</math> angle marked</li> <li>■ The angle on the straight line with <math>h</math> is supplementary with <math>80</math> so <math>180 - 80 = 100</math>, then <math>h = 180 - 100</math></li> <li>■ For the bottom trapezium, <math>h + 60 + 120 + 100 = 360</math>, so <math>h = 360 - 280</math></li> </ul>				<p>✓ <i>Minimally acceptable reason</i> eg</p> <ul style="list-style-type: none"> <li>♦ Alternate</li> <li>♦ Supplementary to <math>80</math>, on a straight line</li> <li>♦ Quadrilateral <math>360 - 280</math></li> </ul> <p>✗ <i>Informal justification without correct geometrical property identified</i> eg</p> <ul style="list-style-type: none"> <li>♦ It's the same as the <math>80</math> because of the parallel lines</li> <li>♦ <math>180 - 100</math></li> <li>♦ <math>360 - 280</math></li> </ul> <p>✗ <i>Incomplete reason</i> eg</p> <ul style="list-style-type: none"> <li>♦ It is the same as the <math>80^\circ</math> angle marked</li> <li>♦ Angles in a quadrilateral add up to <math>360^\circ</math></li> <li>♦ It's opposite the <math>80^\circ</math> on the other side</li> </ul>		
				(U1)							
				1m	<p>Gives <math>j = 120</math> and gives a correct reason eg</p> <ul style="list-style-type: none"> <li>■ The angle on a straight line with <math>j</math> is <math>60</math> because it is an alternate (or corresponding) angle with the <math>60</math> marked, so <math>j = 180 - 60</math></li> <li>■ It's a supplementary angle with angle B so it's <math>180 - 60</math></li> <li>■ For the bottom trapezium, <math>j + 100 + 80 + 60 = 360</math>, so <math>j = 360 - 240</math></li> <li>■ In the parallelogram, angles A and C are equal, so <math>j = (360 - 60 - 60) \div 2</math></li> <li>■ Angle C is supplementary with the <math>60^\circ</math> marked so is <math>180 - 60 = 120</math> <math>j</math> is the opposite angle in the parallelogram to angle C</li> </ul>				<p>✓ <i>Minimally acceptable reason</i> eg</p> <ul style="list-style-type: none"> <li>♦ Alternate (or corresponding), on a straight line</li> <li>♦ Supplementary to <math>60</math></li> <li>♦ Quadrilateral <math>360 - 240</math></li> <li>♦ Parallelogram <math>240 \div 2</math></li> <li>♦ Parallelogram <math>180 - B</math></li> </ul> <p>! <i>For angle j, follow through</i> Accept as <math>200 -</math> their <math>h</math>, alongside a correct reason referring to the quadrilateral containing both angles</p> <p>✗ <i>Informal justification without correct geometrical property identified</i> eg</p> <ul style="list-style-type: none"> <li>♦ <math>180 - 60</math></li> <li>♦ <math>360 - 240</math></li> <li>♦ <math>240 \div 2</math></li> <li>♦ <math>180 - B</math></li> </ul> <p>✗ <i>Incomplete reason</i> eg</p> <ul style="list-style-type: none"> <li>♦ It is the same as angle C which is <math>120^\circ</math></li> <li>♦ Angles in a quadrilateral add up to <math>360^\circ</math></li> <li>♦ <math>j</math> and <math>60</math> are angles on a straight line so add up to <math>180^\circ</math></li> </ul>		
				(U1)							



Tier & Question				Area	
3-5	4-6	5-7	6-8		
		25	17		
				Correct response	Additional guidance
			2m	$100 - \frac{25\pi}{2}$ or 60.7(...) or 60.8 or 61	<p>✓ <i>Pupil working in mm<sup>2</sup></i>                      For 2m, accept values in the correct response column × 100                      For 1m, accept values or methods in the correct response column × 100</p> <p>! <i>The only error is to use the area of a whole circle rather than half a circle</i>                      eg                      ♦ <math>100 - 25\pi</math>                      ♦ 21.4(...) or 21.5 or 21                      Mark as 1, 0</p> <p>✗ <i>Conceptual error</i>                      eg                      ♦ <math>10^2 - 5^2 \times \pi \div 2 = 20 - 5\pi</math>                      ♦ <math>100 - 2 \times \pi \times 5 = 68.6</math></p>
			or 1m	Shows the value $\frac{25\pi}{2}$ or 39.(...), or the value $\frac{25\pi}{4}$ or 19.6(...)  or  Shows a complete correct method with not more than one computational or rounding error eg ■ $10^2 - 5^2 \times \pi \div 2$ ■ $25 \times \pi \div 2 = 40$ (rounding error), $100 - 40 = 60$	
				(U1)	
			1m	Shows the correct unit for their area or method eg ■ 60.8 cm <sup>2</sup> ■ 39.(...) and cm <sup>2</sup> seen ■ 100 and cm <sup>2</sup> seen ■ 6073 mm <sup>2</sup> ■ $100^2 - 50^2 \times \pi \div 2$ and mm <sup>2</sup> seen	<p>! <i>Incorrect or no working or value for area seen, but units given</i>                      If neither mark for calculating the shaded area has been awarded, condone cm<sup>2</sup> seen for the final mark</p>

**Fir trees**

Tier & Question				18	Correct response	Additional guidance																									
3-5	4-6	5-7	6-8																												
				3m	<p>Gives a correct cost of £3332 to £3348 inclusive, and shows or implies a correct method for their cost</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ <math>21 \text{ [value A]} \times 18 = 378</math>  <math>(119 - 21 \text{ [value A]}) \times 22 = 98 \text{ [value B]} \times 22 = 2156</math>  <math>(150 - 119) \times 26 = 31 \text{ [value C]} \times 26 = 806</math>  <math>378 + 2156 + 806 = \text{£}3340</math></li> <li>■ <math>20 \text{ [value A]} \times 18 = 360</math>  <math>100 \text{ [value B]} \times 22 = 2200</math>  <math>30 \text{ [value C]} \times 26 = 780</math>                      Answer £3340</li> <li>■ <math>360 + 2200 + 780 = 3340</math></li> </ul>	<p>Note to markers: For the number of trees in each height range, accept values within the following ranges:</p> <p>Value A: <math>1.2\text{m} &lt; h \leq 1.5\text{m}</math> 20 to 22 inclusive [accurate value 21]</p> <p>Value B: <math>1.5\text{m} &lt; h \leq 1.75\text{m}</math> 118 to 120 inclusive – their A [accurate value 98]</p> <p>Value C: <math>1.75\text{m} &lt; h \leq 2\text{m}</math> 150 – their B – their A [accurate value 31]</p> <p>Note that correct values <b>must</b> follow through</p> <p>Markers may find the following totals useful:</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th colspan="2"></th> <th colspan="3">1<sup>st</sup> reading</th> </tr> <tr> <th colspan="2"></th> <th>20</th> <th>21</th> <th>22</th> </tr> </thead> <tbody> <tr> <td></td> <td>118</td> <td>3348</td> <td>3344</td> <td>3340</td> </tr> <tr> <td>2<sup>nd</sup> reading</td> <td>119</td> <td>3344</td> <td>3340</td> <td>3336</td> </tr> <tr> <td></td> <td>120</td> <td>3340</td> <td>3336</td> <td>3332</td> </tr> </tbody> </table>			1 <sup>st</sup> reading					20	21	22		118	3348	3344	3340	2 <sup>nd</sup> reading	119	3344	3340	3336		120	3340	3336	3332
		1 <sup>st</sup> reading																													
		20	21	22																											
	118	3348	3344	3340																											
2 <sup>nd</sup> reading	119	3344	3340	3336																											
	120	3340	3336	3332																											
				or 2m	<p>Shows a complete correct method with not more than one error</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ <math>21 \times 18 = 378</math>  <math>89 \text{ (error)} \times 22 = 1958</math>  <math>40 \times 26 = 1040</math>                      Answer £3376</li> </ul> <p>or</p> <p>Shows the values 20 to 23 inclusive [value A], 117 to 120 inclusive – their A [value B] and 150 – their B – their A [value C]</p>																										
				or 1m	<p>Shows the values 20 to 23 inclusive, 117 to 120 inclusive and 150</p> <p>or</p> <p>Shows a complete correct method with not more than two errors</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ <math>24 \text{ (error)} \times 18 = 432</math>  <math>100 \text{ (error)} \times 22 = 2200</math>  <math>26 \times 26 = 676</math>                      Answer £3308</li> </ul>	<p>✗ For 1m, values obtained by dividing 150, not reading from the graph</p> <p>eg</p> <ul style="list-style-type: none"> <li>◆ <math>150 \div 3 = 50,</math>  <math>50 \times 18 = 900</math>  <math>50 \times 22 = 1100</math>  <math>50 \times 26 = 1300</math>                      Answer £3300</li> </ul>																									

Tier & Question									<b>Changing shape</b>	
3-5	4-6	5-7	6-8							
				<b>19</b>			<b>Correct response</b>			<b>Additional guidance</b>
			a	2m	21					
				or 1m	Shows a correct method eg <ul style="list-style-type: none"> <li>■ <math>(1.1)^2</math></li> <li>■ Digits 121 seen</li> </ul>					<p><b>!</b> <i>Method uses a numerical value for the sides of the square</i></p> <p>For 1m, accept a complete correct method with not more than one computational error eg, for a square of side 6</p> <ul style="list-style-type: none"> <li>◆ <math>6.6^2 \div 36 \times 100 = 124</math> (error)</li> </ul> <p>Answer: 24%</p> <p>Do not accept a conceptual error such as doubling rather than squaring, or any other error that would lead to a percentage decrease rather than a percentage increase</p>
			b	2m	4 (decrease) or -4					<p>✓ For 2m, 4 with no indication of 'decrease'</p> <p>✗ For 2m, indication of a 4% increase</p>
				or 1m	Indicates a 4% increase  or  Shows or implies a complete correct method with not more than one error eg <ul style="list-style-type: none"> <li>■ <math>100 - \frac{120 \times 80}{100}</math></li> <li>■ Digits 96 seen, with no evidence of an incorrect method</li> <li>■ <math>1.2 \times 0.8 = 0.92</math> (error), so 8%</li> <li>■ 20% of 100 = 20, 100 + 20 = 120, 20% of 120 = 26 (error), 120 - 26 = 94, so 6%</li> </ul>					<p><b>!</b> <i>Method uses numerical values for the sides of the rectangle</i></p> <p>Mark as for part (a) but note that there must be a percentage decrease rather than a percentage increase</p>



Tier & Question				<b>Which graph?</b>		
3-5	4-6	5-7	6-8			
			<b>20</b>		<b>Correct response</b>	<b>Additional guidance</b>
			a	1m	Indicates graph D	
			b	1m	Indicates graph C	
			c	1m	Indicates graph B	

Tier & Question								<b>Side and angle</b>	
3-5	4-6	5-7	6-8						
			<b>21</b>					<b>Correct response</b>	<b>Additional guidance</b>
			a	2m	17 or 17.2(...), with no evidence of accurate or scale drawing				
				<i>or</i> 1m	Shows or implies a correct method with not more than one computational or rounding error eg <ul style="list-style-type: none"> <li>■ <math>28 \times \cos 52</math></li> <li>■ <math>\cos 52 = 0.62</math> (<i>premature rounding</i>), <math>28 \times 0.62 = 17.36</math></li> <li>■ <math>28\sin 38</math></li> </ul> or Shows a correct trigonometric ratio eg <ul style="list-style-type: none"> <li>■ <math>\cos 52 = \frac{w}{28}</math></li> <li>■ <math>\sin 38 = \frac{w}{28}</math></li> </ul>			<p><b>!</b> <i>For 1m, incomplete notation that omits the angle</i> eg  <ul style="list-style-type: none"> <li>◆ <math>\cos = \frac{w}{28}</math></li> </ul>                     Do not accept unless evaluation or other indication shows that the relevance of the angle has been understood</p>	
			b	2m	35 or 34.9(...), with no evidence of accurate or scale drawing				
				<i>or</i> 1m	Shows or implies a complete correct method with not more than one computational or rounding error eg <ul style="list-style-type: none"> <li>■ <math>\tan^{-1} \frac{42}{60}</math></li> <li>■ <math>\tan^{-1} 0.7</math></li> <li>■ Answer of 34</li> </ul> or Shows a correct trigonometric ratio eg <ul style="list-style-type: none"> <li>■ <math>\tan x = \frac{42}{60}</math></li> <li>■ <math>\tan y = \frac{60}{42}</math> [unmarked angle labelled as y]</li> </ul> or The only error is to find the unmarked angle, ie gives an answer of 55 or 55.1(...), with no evidence of accurate or scale drawing			<p><b>✓</b> <i>For 1m, incomplete but unambiguous notation</i> eg  <ul style="list-style-type: none"> <li>◆ <math>\tan = \frac{42}{60}</math></li> </ul> </p>	

Tier & Question					22	Correct response	Additional guidance
3-5	4-6	5-7	6-8				
			a	1m	Shows or implies correct substitution into the formula with correct evaluation of at least the part in brackets eg <ul style="list-style-type: none"> <li>■ Value between 1134 and 1147 inclusive</li> <li>■ 1150</li> <li>■ <math>365\pi</math></li> <li>■ <math>\frac{1}{3} \times \pi \times 5 \times 219</math></li> <li>■ <math>5.2(\dots) \times 219</math></li> </ul>	<p><b>!</b> <i>For the first mark, value(s) rounded</i></p> <p>For <math>\frac{1}{3}</math>, accept 0.33 or better</p> <p>For <math>\pi</math>, accept 3.14 or 3.142 or better</p> <p>eg, for the first mark accept</p> <ul style="list-style-type: none"> <li>♦ <math>0.33 \times 3.14 \times 5 \times 219</math></li> <li>♦ <math>5.1(\dots) \times 219</math></li> </ul>	
				1m	Shows the correct value for the volume of the bowl to 3 significant figures, ie 1150	<p><b>!</b> <i>For the second mark, follow through from an incorrect volume or incorrect working</i></p> <p>Accept provided their volume is greater than 1000, and needs rounding to be given correct to 3 significant figures</p> <p>eg, from their volume as 1031.(...) or working of <math>4.71(\dots) \times 219</math> accept</p> <ul style="list-style-type: none"> <li>♦ 1030</li> </ul> <p>eg, from their volume as 1030 with no working, do not accept</p> <ul style="list-style-type: none"> <li>♦ 1030</li> </ul>	
			b	1m	Gives a correct formula eg <ul style="list-style-type: none"> <li>■ <math>\frac{1}{3} \pi a^2 h</math></li> <li>■ <math>\frac{\pi h a^2}{3}</math></li> </ul>	<p><b>!</b> <i>Unconventional notation</i></p> <p>Condone</p> <p>eg, accept</p> <ul style="list-style-type: none"> <li>♦ <math>\pi \times h \times a \times a \div 3</math></li> </ul> <p><b>✗</b> <i>Formula not completely simplified</i></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ <math>\frac{\pi h a^3}{3a}</math></li> </ul> <p><b>✗</b> <i>Incorrect name for variable within the context of the question</i></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ <math>\frac{1}{3} \pi r^2 h</math></li> </ul>	

Tier & Question								<b>Two circles</b>	
3-5	4-6	5-7	6-8						
			<b>23</b>			<b>Correct response</b>		<b>Additional guidance</b>	
		a	1m			<p>Gives a correct explanation</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ Since BC is a diameter of the smaller circle, any angle made by joining points B and C to a point on the circle's circumference must be <math>90^\circ</math></li> <li>■ BC is a diameter (given) A is on the circumference (intersection of circles)</li> <li>∴ <math>\angle BAC = 90</math></li> <li>■ Angle BAC is an angle in a semicircle, so it must be a right angle</li> </ul>		<p>✓ <i>Minimally acceptable explanation</i></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ BC is a diameter</li> <li>♦ Angles in a semicircle</li> </ul> <p>✗ <i>Incomplete or incorrect explanation</i></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ Angle BAC must be <math>90^\circ</math></li> <li>♦ Semicircle</li> <li>♦ AB is a radius of the large circle, and AC is a tangent of the larger circle, so they must be at right angles</li> </ul>	
		b	2m			8, with no evidence of accurate or scale drawing			
			<i>or</i> 1m			<p>Shows the value 64</p> <p>or</p> <p>Shows sufficient working to indicate correct application of Pythagoras' theorem</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ <math>10^2 - 6^2</math></li> <li>■ <math>\sqrt{100 - 36}</math></li> <li>■ <math>10 \times 10 - 6 \times 6</math></li> </ul> <p>or</p> <p>States or implies that triangle ABC is an enlargement of a 3, 4, 5 right-angled triangle</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ It's a 3, 4, 5 triangle with sides <math>\times 2</math></li> </ul> <p>or</p> <p>Shows a complete correct method with not more than one computational error</p> <p>eg</p> <ul style="list-style-type: none"> <li>■ <math>AC^2 = 11^2</math> (error) <math>- 6^2</math> = 85 AC = 9.2</li> </ul>		<p>✗ <i>For 1m, error is to square then add rather than subtract</i></p> <p>eg</p> <ul style="list-style-type: none"> <li>♦ <math>AC^2 = 10^2 + 6^2</math></li> </ul>	

## Index to mark schemes

Tier				Question	Page
3-5	4-6	5-7	6-8		
1				4 by 4 grid	11
2				Heating	12
3				Tickets	13
4				Unit	14
5				Paralympics	15
6				Half price	16
7				Teachers	16
8	1			Membership	16
9	2			Factor	17
10	3			Shapes on a grid	18
11	4			Meal	18
12	5			Rhombus area	19
13	6			Mobile phones	20
14	7			Arranging numbers	20
15	8			What shape?	21
17	9	1		Algebra grids	22
16	10	2		1976 v 2002	23
18	11	3		Pens	24
20	12	4		Counters	25
19	13	5		From London	25
21	14	6		How many?	26
22	15	7		Pentagon	27
23	16	8		Using a calculator	27
	17	9	1	Tennis prizes	28
	18	10	2	Enlargement	29
	19	11	3	Heron of Alexandria	30
	20	12	4	Hands	31
	21	13	5	Screens	31
	22	14	6	Spinning	32
	23	15	7	Number	32
	24	16	8	A level results	33
	25	17	9	Solutions	34
	26	18	10	Simplify	36
		19	12	Watching	37
		20	11	Milk	38
		21	13	Sequences	40

Tier				Question	Page
3-5	4-6	5-7	6-8		
		22	14	Bracket multiplication	40
		23	15	Parallelogram	41
		24	16	Rich and poor	42
		25	17	Area	43
			18	Fir trees	44
			19	Changing shape	45
			20	Which graph?	46
			21	Side and angle	47
			22	Bowl	48
			23	Two circles	49

First published in 2005

© Qualifications and Curriculum Authority 2005

Reproduction, storage, adaptation or translation, in any form or by any means, of this publication is prohibited without prior written permission of the publisher, unless within the terms of licences issued by the Copyright Licensing Agency. Excerpts may be reproduced for the purpose of research, private study, criticism or review, or by educational institutions solely for educational purposes, without permission, provided full acknowledgement is given.

Produced in Great Britain by the Qualifications and Curriculum Authority under the authority and superintendence of the Controller of Her Majesty's Stationery Office and Queen's Printer of Acts of Parliament.

The Qualifications and Curriculum Authority is an exempt charity under Schedule 2 of the Charities Act 1993.

Qualifications and Curriculum Authority  
83 Piccadilly  
London  
W1J 8QA  
[www.qca.org.uk/](http://www.qca.org.uk/)

***Further teacher packs may be purchased (for any purpose other than statutory assessment) by contacting:***

QCA Orderline, PO Box 29, Norwich NR3 1GN  
tel: 08700 606015; fax: 08700 606017  
email: [orderline@qca.org.uk](mailto:orderline@qca.org.uk)

**Order ref:** QCA/05/1427

265512